

The importance of N_2 leptogenesis

Guy Engelhard,¹ Yuval Grossman,² Enrico Nardi,^{3,4} and Yosef Nir¹

¹*Department of Particle Physics, Weizmann Institute of Science, Rehovot 76100, Israel*

²*Department of Physics, Technion-Israel Institute of Technology, Technion City, Haifa 32000, Israel*

³*INFN, Laboratori Nazionali di Frascati, C.P. 13, 100044 Frascati, Italy*

⁴*Instituto de Física, Universidad de Antioquia, A.A.1226, Medellín, Colombia*

(Dated: February 2, 2008)

We argue that fast interactions of the lightest singlet neutrino N_1 would project part of a preexisting lepton asymmetry L_p onto a direction that is protected from N_1 washout effects, thus preventing it from being erased. In particular, we consider an asymmetry generated in N_2 decays, assuming that N_1 interactions are fast enough to bring N_1 into full thermal equilibrium. If N_1 decays occur at $T \gtrsim 10^9$ GeV, that is, before the muon Yukawa interactions enter into thermal equilibrium, then generically part of L_p survives. In this case some of the constraints implied by the standard N_1 leptogenesis scenario hold only if $L_p \approx 0$. For $T \lesssim 10^9$ GeV, L_p is generally erased, unless special alignment/orthogonality conditions in flavor space are realized.

Introduction. The existence of heavy singlet neutrinos with Majorana masses is quite generic in theories that go beyond the Standard Model (SM) and, furthermore, can simultaneously explain two puzzles: It can induce, via the see-saw mechanism, active neutrino masses that are much lighter than those of the SM charged fermions, and it can generate, via leptogenesis [1], the baryon asymmetry of the Universe (BAU).

Leptogenesis is of major phenomenological interest because, within the conventional leptogenesis scenario, the measured value of the BAU has non-trivial implications for other fundamental issues. In particular, it provides constraints on low energy neutrino parameters, on the mass scale of the heavy Majorana neutrinos, and on the reheating temperature after inflation. In order to understand how robust are such implications, it is important to explore under which conditions the conventional picture holds. The aim of this paper is to show that, even within the standard scenario, one of the assumptions that underlies most of the leptogenesis analyses might not hold, implying that the standard leptogenesis constraints are relaxed or even evaded.

In the conventional picture, three singlet neutrinos N_α are added to the SM, with hierarchical Majorana masses $M_1 \ll M_2 \ll M_3$. It is often assumed that the L -violating interactions of N_1 would washout any lepton asymmetry L_p generated at temperatures $T \gg M_1$ and, in particular, the asymmetry generated in the decays of $N_{2,3}$. If this were the case, the final asymmetry would depend only on N_1 dynamics, and the number of parameters relevant for leptogenesis would be reduced to just a few: the mass of the lightest singlet neutrino M_1 , the “washout parameter” $\tilde{m}_1$ (defined below), and the CP asymmetry in N_1 decays ϵ_{N_1} . We show that, contrary to common wisdom, part of the lepton asymmetry generated in $N_{2,3}$ decays in general survives the N_1 lepto-

genesis phase. Thus, it is quite possible that the lepton asymmetry relevant for baryogenesis originates mainly from $N_{2,3}$ decays. The possibility that N_2 leptogenesis could successfully explain the BAU has been already suggested in some recent papers [2, 3, 4]. However, the scenario put forth in these analyses is essentially that of “ N_1 -decoupling”, in which the Yukawa couplings of N_1 are simply too weak to washout the N_2 -generated asymmetry. Here we point out that N_1 decoupling is not a necessary condition, and that N_2 leptogenesis can be successful even when the Yukawa interactions of N_1 are fast enough to bring these states into complete thermal equilibrium. Our main conclusions, regarding the survival of part of the asymmetry generated in N_2 decays, have been previously stated in refs. [5, 6]. We describe the framework where this picture arises, spelling out the assumptions involved and uncovering some important subtleties, and we explain the importance of the results for the analysis of leptogenesis.

Notations. The particles that play a role in leptogenesis are the heavy singlet neutrinos N_α , the light lepton $SU(2)$ -doublets L_i and $SU(2)$ -singlets E_i , and the standard model Higgs H . The relevant Lagrangian terms are

$$-\mathcal{L} = \frac{1}{2}M_\alpha N_\alpha N_\alpha + \lambda_{\alpha i} H N_\alpha L_i + Y_i H^\dagger L_i E_i. \quad (1)$$

where $\alpha = 1, 2, 3$ is a heavy neutrino index while $i = e, \mu, \tau$ is a flavor index. Eq. (1) is written in the mass basis for the singlet neutrinos and for the charged leptons, that is, M and Y are diagonal. It is convenient to define ℓ_α , the three (in general non-orthogonal) combinations of lepton doublets to which the corresponding N_α decay:

$$|\ell_\alpha\rangle = (\lambda\lambda^\dagger)^{-1/2}_{\alpha\alpha} \sum_i \lambda_{\alpha i} |L_i\rangle. \quad (2)$$

We denote by n_{ℓ_α} the number density of ℓ_α and normalize

the asymmetry-density to the equilibrium density n_{eq} :

$$y_{\ell_\alpha} = (n_{\ell_\alpha} - n_{\bar{\ell}_\alpha})/n_{\text{eq}}. \quad (3)$$

We use similar notations for other particle species, *e.g.* n_H and y_H for the Higgs.

The CP asymmetry generated in N_α decays reads:

$$\epsilon_{N_\alpha} = \frac{\Gamma(N_\alpha \rightarrow \ell H) - \Gamma(N_\alpha \rightarrow \bar{\ell} \bar{H})}{\Gamma(N_\alpha \rightarrow \ell H) + \Gamma(N_\alpha \rightarrow \bar{\ell} \bar{H})}. \quad (4)$$

We introduce three quantities $\tilde{m}_\alpha$ with dimension of a mass that parametrize the washout of the lepton asymmetry ϵ_{N_α} due to N_α -related interactions:

$$\tilde{m}_\alpha = \frac{\langle H \rangle^2 (\lambda \lambda^\dagger)_{\alpha\alpha}}{M_\alpha}. \quad (5)$$

The baryon number generated from the decays of the $N_{1,2,3}$ neutrinos, normalized to the entropy density s can be formally written as follows:

$$Y_B \equiv \frac{n_B - n_{\bar{B}}}{s} = -1.38 \times 10^{-3} \sum_{\alpha,\beta} \epsilon_{N_\alpha} \eta_{\alpha\beta}, \quad (6)$$

where $\eta_{\alpha\beta}$ are efficiency factors that are related to the effects of N_β interactions on the asymmetry ϵ_{N_α} [7]. The diagonal factors $\eta_{\alpha\alpha}$ are simply related to washout effects. The off-diagonal factors $\eta_{\alpha\beta}$ ($\alpha \neq \beta$) have a more complicated structure since, as we discuss below, they represent also decoherence effects on ℓ_α induced by N_β interactions. In the standard leptogenesis scenario (and in the one-flavor approximation), only one efficiency factor is relevant, η_{11} . Calculating this factor requires solving the complete set of Boltzmann equations. However, for $M_1 \ll 10^{14}$ GeV (when $\Delta L = 2$ washouts can be neglected) η_{11} is completely determined by the mass and couplings of N_1 via the combination $\tilde{m}_1$ defined in Eq. (5). A simple relation that approximates quite well the dependence of η_{11} on $\tilde{m}_1$ can be found *e.g.* in [9].

Recent data on the cosmic background anisotropy [10] and considerations of Big Bang Nucleosynthesis [11] yield

$$Y_B^{\text{exp}} = (8.7 \pm 0.3) \times 10^{-11}. \quad (7)$$

This is the experimental number that leptogenesis aims to explain. In the following we discuss the possibility that this number could be related to ϵ_{N_2} , rather than to ϵ_{N_1} .

The framework. We are interested in the strong washout regime for N_1 -related interactions:

$$\tilde{m}_1 \gg m_*, \quad (8)$$

where $m_* = 1.66\sqrt{g_*}16\pi\langle H \rangle^2/M_P \approx 10^{-3}$ eV, g_* is the number of degrees of freedom (d.o.f.) in equilibrium and M_P is the Planck mass. In the weak washout regime ($\tilde{m}_1 \lesssim m_*$) and in the N_1 decoupling regime ($\tilde{m}_1 \ll m_*$) the final $B - L$ asymmetry depends in general on the

initial conditions for N_1 leptogenesis. We will find that, contrary to common belief, this is true also for the regime of strong N_1 washouts.

We consider a lepton asymmetry $L_p \neq 0$ that is present before the onset of N_1 leptogenesis. For definiteness we study the case where L_p originates from out-of-equilibrium, CP violating decays of N_2 . (The effects of N_3 decays can be readily included in the same way as N_2 effects, and should be taken into account in quantitative estimates of the BAU resulting from leptogenesis.) We assume that N_2 -related washouts are not too strong:

$$\tilde{m}_2 \not\gg m_*, \quad (9)$$

so that a finite L_p survives [12].

Our analysis is qualitative. To simplify our arguments, we impose two further conditions: thermal leptogenesis, that is $n_{N_2}(T \gg M_2) \approx 0$ and in particular

$$n_{N_1}(T \gg M_1) \approx 0, \quad (10)$$

and strong hierarchy,

$$M_2/M_1 \gg 1. \quad (11)$$

We now describe the simplifications that occur when the conditions (8)-(11) are realized. The hierarchy (11) is assumed to be large enough that, together with Eq. (10), $n_{N_1}(T \sim M_2) \approx 0$ is guaranteed. On the other hand, the Boltzmann suppression gives $n_{N_2}(T \sim M_1) \approx 0$. This situation has several consequences: (a) there are no N_1 related washout effects during the N_2 leptogenesis, (b) there are no N_2 related washout effects during N_1 leptogenesis, (c) we do not need a density matrix to describe the ℓ_1 and ℓ_2 densities during N_2 leptogenesis, and (d) the density matrix for ℓ_1 and for the states orthogonal to ℓ_1 does not depend on the interactions with N_2 .

Condition (8) implies that, already at $T \gtrsim M_1$, the interactions mediated by the N_1 Yukawa couplings become sufficiently fast to quickly destroy the coherence of the state ℓ_2 produced in N_2 decays. Then a statistical mixture of ℓ_1 and of the states orthogonal to ℓ_1 builds up, and it can be described by a *diagonal* density matrix. The analogous situation in which decoherence occurs because of fast charged lepton Yukawa interactions was analyzed in [5, 8, 13, 14, 15] in the context of flavored leptogenesis. It was found [14, 15] that after a short transient period during which the flavor-components of the asymmetry undergo fast oscillations, they decohere and get projected onto the flavor basis. There is, however, a difference between the present case and the flavor case: as the temperature drops below M_1 , the N_1 decohering interactions are suppressed by T^2/M_1^2 and thus slow down with respect to the Universe expansion rate. When $T/M_1 < m_*/\tilde{m}_1$, they become irrelevant. Our assumption (8) implies that this happens only at $T \ll M_1$, that is after quantum coherence between the ℓ_2 components has been completely

destroyed. Finally, it is crucial that at $T \sim M_1$ the effects related to N_2 interactions are suppressed enough not to interfere with the decoherence effects of N_1 , and this is ensured if the condition $M_1/M_2 < m_*/\tilde{m}_2$ holds. The combination of Eq. (9) with Eq. (11) implies that this condition is indeed fulfilled.

On general grounds one would expect that decoherence effects proceed faster than washout. In the relevant range, $T \gtrsim M_1$, this is also ensured by the fact that the dominant $\mathcal{O}(\lambda^2)$ washout processes, that is the inverse decays $\ell H \rightarrow N_1$, are blocked because of thermal effects [9], and only scatterings with top-quarks and gauge bosons, that have additional suppression factors of Y_t^2 and g^2 , contribute to the washout.

In the following we analyze different temperature regimes, paying attention to the cases where the interplay between flavor effects and decoherence effects induced by the N_1 Yukawa couplings becomes important.

$M_{1,2} \gtrsim 10^{12}$ GeV. We first consider the case where both N_2 and N_1 decay at $T \gtrsim 10^{12}$ GeV, *i.e.* when all charged lepton Yukawa interactions are slower than the expansion rate of the Universe and flavor effects are irrelevant. (Note that $\Delta L = 2$ interactions of the heavier neutrinos can generally be in equilibrium down to $T \sim 10^{13}$ GeV. Taking these interactions into account complicates the analysis considerably. We thus neglect these interactions, keeping in mind that the range of temperatures and parameters where our analysis applies straightforwardly is rather restricted.)

During the stage of N_2 decays we have, as explained above, $n_{N_1} \approx 0$. Consequently, there are no N_1 -related washout effects during N_2 leptogenesis. Since we assume $\eta_{22} \not\ll 1$, sizeable asymmetries can survive the N_2 -related washouts and, since we neglect N_3 -related effects, we denote the total lepton asymmetry (corresponding to the trace of density matrix of the asymmetries) present at the end of N_2 leptogenesis by y_{ℓ_2} . We thus have

$$y_{\ell_2} \sim \eta_{22}\epsilon_{N_2}. \quad (12)$$

Decoherence effects induced by the N_1 Yukawa interactions become important at a later stage. For $\tilde{m}_1$ large enough (8), they become faster than the expansion rate of the Universe already at $T \gtrsim M_1$. To study the fate of y_{ℓ_2} through the N_1 leptogenesis phase, it is convenient to choose the (orthogonal) basis $(\ell_1, \ell_0, \ell'_0)$ where, without loss of generality, $\langle \ell'_0 | \ell_2 \rangle = 0$. Due to the invariance of the trace under a change of basis, the lepton asymmetry y_{ℓ_2} of Eq. (12) decomposes into the two components:

$$y_{\ell_2} = y_{\ell_0} + y_{\ell_1}. \quad (13)$$

In general (and, in particular, if there is no fine-tuned alignment of ℓ_1 and ℓ_2) we expect $y_{\ell_0} = \mathcal{O}(y_{\ell_2})$. The crucial point is that since ℓ_0 is orthogonal to ℓ_1 , y_{ℓ_0} is protected against N_1 washouts. This ensures that a finite

part of the asymmetry generated in N_2 decays survives the N_1 leptogenesis phase, and is a source of $Y_L \neq 0$.

Actually, by taking into account the constraints on the various chemical potentials implied by the reactions that are in equilibrium in this temperature range, one can get a better insight about the true composition of the surviving lepton asymmetry. To simplify the analysis let us assume that $\tilde{m}_1$ is so large that around $T \sim M_1$, when the number density of N_1 is maximal, N_1 -interactions attain chemical equilibrium. This implies:

$$y_{N_1} = y_{\ell_1} + y_H/2, \quad (14)$$

where y_{N_1} corresponds to a chemical potential for N_1 that we will eventually set to zero. Combining (14) with the conditions implied by in-equilibrium gauge and top-Yukawa interactions and by baryon number and hypercharge conservation, we obtain

$$\begin{aligned} y_H &= (2/3)(y_{\ell_1} + y_{\ell_0}), \\ y_{N_1} &= (4/3)y_{\ell_1} + (1/3)y_{\ell_0}, \\ Y_L/Y^{\text{eq}} &= (10/3)y_{\ell_1} + (7/3)y_{\ell_0}, \end{aligned} \quad (15)$$

where Y^{eq} is the equilibrium density of relativistic species with one d.o.f.. At temperatures $T \sim M_1$ the L -violating interactions of N_1 are fast and imply $y_{N_1} \rightarrow 0$, finally yielding $y_H = y_{\ell_0}/2$ and $y_{\ell_1} = -y_{\ell_0}/4$. We note that since the strong washout (equilibrium) condition (14) does not imply $y_{\ell_1} = 0$ and $y_H = 0$ separately, part of the asymmetry in ℓ_1 , that is a quarter in size and opposite in sign to y_{ℓ_0} , also survives, yielding a total lepton asymmetry that is somewhat smaller than $2y_{\ell_0}$:

$$Y_L/Y^{\text{eq}} = (3/2)y_{\ell_0}. \quad (16)$$

In a fine-tuned situation where $\ell_2 \parallel \ell_1$ and $y_{\ell_0} = 0$, y_{ℓ_2} can be completely washed out by N_1 interactions. (This alignment condition would entail one massless neutrino.) In this case, however, some fraction of y_{ℓ_3} generated in N_3 decays remains protected from $N_{1,2}$ washouts. One cannot further impose $\ell_3 \parallel \ell_2 \parallel \ell_1$ since this would imply two massless neutrinos.

10^9 GeV $\lesssim M_1 \lesssim 10^{12}$ GeV $\lesssim M_2$. We now consider the case where (a) N_1 decays at 10^9 GeV $\lesssim T \lesssim 10^{12}$ GeV, when the τ (but not the μ) Yukawa interactions are in equilibrium, while (b) N_2 leptogenesis occurs in the “flavor blind” regime, $T \gtrsim 10^{12}$ GeV yielding the result in Eq. (12). To study the onset of fast N_1 interactions, we use the orthogonal basis $(\ell_\tau, \ell_a, \ell_0)$ where, without loss of generality, $\langle \ell_0 | \ell_1 \rangle = 0$ while (except for the specific cases where ℓ_τ is either aligned with or orthogonal to ℓ_1) $\langle \ell_{\tau,a} | \ell_1 \rangle \neq 0$. Given that ℓ_0 is affected by neither τ nor N_1 interactions, its off-diagonal quantum correlations are quickly damped, singling out the ℓ_0 component of ℓ_2 . The crucial quantity is then y_{ℓ_0} , the projection of the lepton asymmetry onto the ℓ_0 direction. We can again conclude that this part of the lepton asymmetry remains protected against N_1 washouts.

Compared to the previous case, however, we now have a larger number of interactions that are in equilibrium. As a representative set for this temperature interval, we take the interactions of the third family Yukawas $Y_{t,b,\tau}$ and the QCD and electroweak sphalerons. The latter violate baryon number, but conserve the three charges $\Delta_i \equiv B/3 - L_i$ (for the case at hand $i = \tau, a, 0$). Furthermore, any initial asymmetry in Δ_0 generated during N_2 decays is conserved also by N_1 interactions. After solving for the equilibrium constraints and setting $y_{N_1} = 0$ we find:

$$Y_{B-L} = +(90/107) Y_{\Delta_0}, \quad (17)$$

$$Y_{\Delta_\tau} = -(16/107) Y_{\Delta_0}, \quad (18)$$

$$Y_{\Delta_a} = -(1/107) Y_{\Delta_0}, \quad (19)$$

where $Y_{\Delta_0} \equiv -2y_{\ell_0} Y^{\text{eq}}$. We see that, in spite of having equilibrium reactions with N_1 , fractions of y_{ℓ_τ} and y_{ℓ_a} survive. In addition, the fast τ Yukawa interactions generate y_τ , and the fast sphalerons generate a net baryon number. The overall effect is to cancel a (small) part of the y_{ℓ_0} contribution to the final $B - L$.

In deriving Eqs. (17)-(19) we assume that ℓ_1 has sizable projections on both the $\ell_{\tau,a}$ states, leaving aside the fine tuned possibilities $\ell_1 \perp \ell_\tau(\ell_a)$. Thus we apply the two equilibrium conditions $y_{\ell_\tau} = y_{\ell_a} = -y_H/2$. In the realistic case of large but not unnaturally large N_1 washouts, the condition $y_{\ell_\tau} = y_{\ell_a}$ is not expected to hold, since the projections of y_{ℓ_2} onto these two states, as well as their specific washouts, differ. Solving for the coupled evolution of y_{ℓ_τ} and y_{ℓ_a} is a non-trivial problem, given that their off-diagonal correlations, that are maintained by the N_1 Yukawa interactions, are likely to be damped only at small T , when the N_1 effects are suppressed by T^2/M_1^2 . The detailed evolution of the ℓ_τ - ℓ_a system represents an interesting quantitative problem, but will not change the main qualitative result: Part of the y_{ℓ_2} asymmetry survives the N_1 leptogenesis phase. If also N_2 leptogenesis occurs at $T \lesssim 10^{12}$ GeV, some details of the previous analysis change, but the final conclusion remains. We conclude that if leptogenesis occurs in the theoretically preferred range $T \gtrsim 10^9$ GeV, when the μ Yukawa reactions are still slow, the constraints inferred by requiring successful N_1 leptogenesis hold only under the additional assumption that either ϵ_{N_2} is small enough or N_2 -related washout is strong enough ($\eta_{22} \ll 1$) to drive $y_{\ell_2} \rightarrow 0$.

$M_1 \lesssim 10^9$ GeV. The last possibility is that N_1 decays at $T \lesssim 10^9$ GeV (the N_2 decay temperature is irrelevant here). In this regime, because of the fast τ and μ Yukawa reactions, the full lepton flavor basis ($\ell_e, \ell_\mu, \ell_\tau$) is resolved. In the general situation, where $\langle \ell_i | \ell_1 \rangle \ll 1$ for $i = e, \mu, \tau$, there are no directions in flavor space where an asymmetry remains protected from N_1 -related washouts, and N_2 leptogenesis fails. A preexisting asymmetry can only survive if it is stored in a flavor that is weakly coupled to N_1 .

Conclusions. The effect of N_1 interactions on N_2 leptogenesis is twofold: decoherence of ℓ_2 , and washout of the lepton-asymmetry y_{ℓ_2} generated in N_2 decays. For weak N_1 washouts ($\tilde{m}_1 \ll 10^{-3}$ eV) neither effect is important, and y_{ℓ_2} survives. For N_1 washouts in the strong regime ($\tilde{m}_1 \gg 10^{-3}$ eV), both effects are important. For a generic flavor structure, N_1 -related decoherence effects project part of y_{ℓ_2} onto a flavor direction that is protected against N_1 washout, and this component of the initial asymmetry survives. Previous analyses ignored N_1 -related decoherence effects, and assumed that when N_1 leptogenesis occurs in the strong washout regime the final BAU is independent of initial conditions. We find that this assumption is justified only in the following cases: *i*) Vanishing decay asymmetries and/or efficiency factors for $N_{2,3}$ ($\epsilon_{N_2} \eta_{22} \approx 0$ and $\epsilon_{N_3} \eta_{33} \approx 0$); *ii*) N_1 -related washouts are still significant at $T \lesssim 10^9$ GeV; *iii*) Reheating occurs at a temperature in between M_2 and M_1 . In all other cases the $N_{2,3}$ -related parameters cannot be ignored when calculating the BAU.

Acknowledgments. We thank Michael Dine, Marta Losada, Michael Peskin, Antonio Riotto, Esteban Roulet, and Adam Schwimmer for useful discussions. This project was supported by the Albert Einstein Minerva Center for Theoretical Physics. The work of Y.G. is supported in part by the Israel Science Foundation under Grant No. 378/05. The work of E.N. is supported in part by Colciencias under contract 1115-333-18739. The research of Y.N. is supported by the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities, and by a grant from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel. E.N. and Y.N. thank the organizers of the International Workshop on High Energy Physics in the LHC Era held in Valparaiso-Chile where this work was completed, for their kind invitation and pleasant working atmosphere.

-
- [1] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).
 - [2] O. Vives, Phys. Rev. D **73**, 073006 (2006).
 - [3] P. Di Bari, Nucl. Phys. B **727**, 318 (2005).
 - [4] S. Blanchet and P. Di Bari, JCAP **0606**, 023 (2006).
 - [5] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, Nucl. Phys. B **575**, 61 (2000).
 - [6] A. Strumia, arXiv:hep-ph/0608347.
 - [7] In the regimes when flavor effects are important, a flavor index also enters in Eq. (6): $\epsilon_{N_\alpha} \eta_{\alpha\beta} \rightarrow \sum_i \epsilon_{N_\alpha}^i \eta_{\alpha\beta}^i$ (see e.g. [8] for the definitions of ‘flavored’ ϵ and η .)
 - [8] E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601**, 164 (2006).
 - [9] G. F. Giudice *et al.*, Nucl. Phys. B **685**, 89 (2004).
 - [10] D. N. Spergel *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **170** (2007) 377.
 - [11] G. Steigman, Int. J. Mod. Phys. E **15**, 1 (2006).
 - [12] It is impossible to satisfy Eq. (9) for the case of quasi-degenerate neutrinos. For this (and other) reasons, we

implicitly assume hierarchical light neutrinos masses.

- [13] T. Endoh, T. Morozumi and Z. h. Xiong, Prog. Theor. Phys. **111**, 123 (2004).
- [14] A. Abada *et al.*, JCAP **0604**, 004 (2006).
- [15] A. De Simone and A. Riotto, JCAP **0702**, 005 (2007).
- [16] E. Nardi, Y. Nir, J. Racker and E. Roulet, JHEP **0601**, 068 (2006).